

DEPARTMENT OF MATHEMATICS, GC UNIVERSITY, LAHORE

Admission Test for M.Phil (Mathematics)/Session 2009-11

Max. Time: 90 MinutesMax. Marks: 60

Name of Candidate _____ Father / Guardian Name _____

Form No. _____ Signature of Candidate _____

Note: Please put a tick (✓) mark on the correct answer in each question. Overwriting will not be evaluated.

1. Let A be a $z \times z$ real matrix such that $A \begin{bmatrix} 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$, $A \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ then A =

(i) $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

(iv) $\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$

(ii) $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$

(v) $\begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$

(iii) $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$

2. If A is a matrix with characteristic equation $x^2 - 4x + 1 = 0$ then $\det A =$

(i) 2

(ii) 0

(iii) -2

(iv) 1

(v) 4

3. Let V and W be two proper subspaces of the space \mathbb{R}^4 . Then $V \cap W$ is a subspace of \mathbb{R}^4 of dimension;

(i) one only

(iv) 0, 1, 2 only

(ii) two only

(v) four only

(iii) zero only

4. In the complex plane, the equation $z^2 = |z|^2$ represents

(i) a pair of points

(iv) a line

(ii) a circle

(v) none of these

(iii) half line

5. The function of $f: \mathbb{C} \rightarrow \mathbb{R}$, from the complex set C into the real set R defined by $f(z) = |z|^2$ is:

(i) differentiable for all z

(iv) continuous only at $z = 0$

(ii) discontinuous for all z

(v) none of these

(iii) differentiable only at $z = 0$

(P.T.O)

6. The alternating group A_4 has,

- | | | | |
|-------|--------------------------|------|---------------------------|
| (i) | five elements of order 3 | (iv) | three elements of order 2 |
| (ii) | two elements of order 2 | (v) | none of these |
| (iii) | one element of order 4 | | |

7. The Quaternion group $\{ \langle a, b : a^4 = e = b^4 = (ab)^4, bab^{-1} = a^{-1} \}$ has

- (i) one element of order 2
 (ii) three elements of order 4
 (iii) seven elements of order 4
 (iv) no element of order 2
 (v) four elements of order 4

8. Let a function $f(z)$ be analytic in a simply connected domain D and C be a closed continuous curve in D . then

- | | | | |
|-------|-----------------------------------|------|----------------------|
| (i) | $\int_C f(z) dz = 1$ | (iv) | $\int_D f(z) dz = 0$ |
| (ii) | $\int_C f(z) dz = 0$ | (v) | none of the above |
| (iii) | $\int_C f(z) dz = \int_D f(z) dz$ | | |

9. If ∇ is the vector differential operator and ϕ is any scalar function then, $\nabla(\nabla\phi) =$

- | | | | |
|-------|----------------|------|-----------------------------|
| (i) | 0 | (iv) | $(\nabla\phi) \cdot \nabla$ |
| (ii) | 1 | (v) | none of the above. |
| (iii) | $\nabla^2\phi$ | | |

10. If $f(x, y, z) = xyz$ is a scalar function and $\nabla = i \frac{\partial}{\partial x} + j \frac{\partial}{\partial y} + k \frac{\partial}{\partial z}$ then $\nabla f =$

- | | | | |
|-------|---|------|-----------------|
| (i) | $x\mathbf{i} + y\mathbf{j} + z\mathbf{k}$ | (iv) | $xyz\mathbf{j}$ |
| (ii) | $xy\mathbf{i}$ | (v) | none of these |
| (iii) | $yz\mathbf{i}$ | | |

11. An integrating factor for the differential equation $\frac{-2y}{x} dx + (x^2 y \cos y + 1) dy = 0$ is,

- | | | | |
|-------|----------------|------|-----------------|
| (i) | 1 | (iv) | x^2 |
| (ii) | $-2x$ | (v) | $\frac{1}{x^2}$ |
| (iii) | $\frac{-2}{x}$ | | |

12. If $f(x) = e^x - e^{-x}$, then $[f'(x)]^2 - [f(x)]^2 =$

- | | | | |
|-------|-----------|------|--------|
| (i) | 4 | (iv) | $2e^x$ |
| (ii) | $4e^{2x}$ | (v) | 2 |
| (iii) | $2e^{-x}$ | | |

13. For which value of k , x^k is a solution for the differential equation $x^2y'' - 3xy' + 4y = 0$?

- (i) 4
 (ii) 3
 (iii) 2
 (iv) 1
 (v) none of these.

14. If $f(x) = \int_1^{x^2} \frac{dt}{1+t^3}$ then $f'(2) =$

- (i) $4/65$
 (ii) $\frac{1}{9}$
 (iii) $\ln\left(\frac{65}{2}\right)$
 (iv) $\ln\left(\frac{9}{2}\right)$
 (v) 0.23

15. The radius of curvature of $f(x) = x + \frac{1}{x}$ at the point P (1, 2) is,

- (i) 1
 (ii) $\sqrt{2}$
 (iii) 4
 (iv) 2
 (v) $\frac{1}{2}$

16. For a pure radioactive model, the rate of decaying variation w.r.t. time t of mass M

satisfies the equation $\frac{dM}{dt} = \frac{-M}{10}$. Find M in terms of its initial value M_0 after 20 units of time t as,

- (i) $\frac{1}{2}M_0$
 (ii) $\frac{1}{4}M_0$
 (iii) $\frac{M_0}{2e}$
 (iv) $\frac{M_0}{e}$
 (v) $\frac{M_0}{e^2}$

17. For $x^2z - 2yz^2 + xy = 0$, find $\frac{\partial x}{\partial z}$ at (1, 1, 1) as;

- (i) 0
 (ii) $4/3$
 (iii) -1
 (iv) 1
 (v) none of these

18. The solution set on the number line $(-\infty, \infty)$ of the inequality $\frac{1}{x-2} < \frac{1}{x+3}$ is the set,

- (i) (-3, -2)
 (ii) (-3, 2)
 (iii) (2, 3)
 (iv) (-2, 2)
 (v) (0, 2)

19. If A is a countable subset of the unit interval $[0, 1]$ then the Lebesgue measure of A is,

- (i) $\frac{1}{2}$
 (ii) 0
 (iii) $2/3$
 (iv) 3^{-1}
 (v) none of these

(P.T.O)

20. The Inequality $\frac{x^2}{a^2} + \frac{y^2}{b^2} < 2$ in \mathbb{R}^2 is,
- (i) a closed set (iv) a dense set
 (ii) an open set (v) none of these
 (iii) a compact set
21. A parabola is homeomorphic to:
- (a) an ellipse (d) a circle
 (b) a straight line (e) none of these
 (c) a hyperbola
22. Which of the following property is NOT true for the interval $[0, 1]$ in \mathbb{R} ?
- (i) it is compact (iv) it is closed
 (ii) it is disconnected (v) none of these
 (iii) it is connected
23. If X is any set, then the collection of all one point subsets of X is a basis for which topology on X :
- (i) confinite (iv) quotient
 (ii) discrete (v) none of these
 (iii) indiscrete
24. Let (X, τ) be a topological space, and $Y \subset X$. Then which of the following defines a subspace topology on Y :
- (i) $\{(Y \cup V)\} : V \in \tau$ (iv) $\{(V - Y)\} : V \in \tau$
 (ii) $\{(Y \cap V)\} : V \in \tau$ (v) none of these
 (iii) $\{(Y - V)\} : V \in \tau$
25. Which of the following is a connected subset of \mathbb{R} ?
- (i) \mathbb{Q} (iv) \mathbb{Z}
 (ii) $[0, 1) \cup (1, 2)$ (v) none of these
 (iii) $[0, \frac{1}{2}) \cup [\frac{1}{2}, 1)$
26. Which of the following is NOT a compact subset of \mathbb{R} ?
- (i) $[0, 1) \cup \{1\}$ (iv) $[2, 3) \cap (3, 4]$
 (ii) $[2, 3) \cup (3, 4]$ (v) none of these
 (iii) $[2, 3]$
27. If $x(y+z) > w$ and $x, y, z,$ and w are all integers, which of the following must be true ?
- (i) $x(y+z) > 0$ (iv) $x+y+z = w$
 (ii) $xy+z=w$ (v) x, y, z and w are all positive
 (iii) $xy+xz=w$

28. In the group \mathbb{Z} of all integers, which of the subset is not its subgroup?

- | | |
|--|---|
| (i) $\{0\}$ | (iv) $\{n \in \mathbb{Z} : n \text{ is even}\}$ |
| (ii) $\{n \in \mathbb{Z} : n > 0\}$ | (v) \mathbb{Z} |
| (iii) $\{m \in \mathbb{Z} : m \text{ is divisible by 6 and 9}\}$ | |

29. Let V_1 and $V_2 (\neq V_1)$ be two 6 dimensional subspaces of a 10-dimensional space V . What is the dimension of $V_1 \cap V_2$?

- | | |
|---------|--------|
| (i) 0 | (iv) 4 |
| (ii) 1 | (v) 6 |
| (iii) 2 | |

30. Let $\mathbb{R}[x]$ be the ring of polynomials in x . Which of the following subsets form its subring;

- | | |
|---|---|
| (i) all the polynomials with coefficient in \mathbb{R} of x is zero | (iii) all the polynomials of odd degree |
| (ii) all the polynomials of even degree | (iv) All the polynomials of degree 2 |
| | (v) none of these |

31. If f is a function defined on the interval $(2,3)$ on the real line $\mathbb{R} = (-\infty, \infty)$ such that $2 < f(x) < 3$, x in $(2,3)$ then;

- | | |
|----------------------------------|---|
| (i) f is bounded | (iv) f is polynomial function of degree 1 |
| (ii) f is negative | (v) f is non-constant |
| (iii) f is strictly increasing | |

32. For what value / values of k , the vector $(1, 2, k, 5)$ of the space \mathbb{R}^4 is a linear combination of the vectors $(0,1,1,1)$, $(0,0,0,1)$ and $(1,1,2,3)$ of \mathbb{R}^4 ;

- | | |
|--------------|---------------------|
| (i) no value | (iv) 3 only |
| (ii) -1 only | (v) infinitely many |
| (iii) 1 only | |

33. The harmonic conjugate $V(x,y)$ for the harmonic function $u(x,y) = y + 3xy^2 - x^3$ is,

- | | |
|-------------------------|-------------------------|
| (i) $-y + 3xy^2 - x^3$ | (iv) $y^3 - 3x^2y + x$ |
| (ii) $x + 3xy^2 - x^2$ | (v) $y^3 - 3x^2y + x^2$ |
| (iii) $y^3 - 3x^2y - x$ | |

34. If A is a 2×2 real matrix then

- | | |
|--|--|
| (i) all the entries of A^2 are none-negative | (iii) if A has two distinct eigen values then so has A^2 |
| (ii) $\det A^2$ is non-negative | (iv) A^2 is a scalar matrix |
| | (v) none of these |

(P.T.O)

35. If S is a non-empty set such that $|S| = k$ and if there are n one-to-one functions on S then $n = ?$
- (i) k^2 (iv) \underline{k}
(ii) k^k (v) none of these
(iii) 2^k
36. If n stands for number of non-isomorphic group of order 4 then $n =$
- (i) 1 (iv) 3
(ii) 0 (v) 5
(iii) 2
37. In the symmetric group S_3 if m stands for the number of subgroup of S_3 then $m = \dots?$
- (i) 1 (iv) 4
(ii) 2 (v) 6
(iii) 3
38. A group of order 144 has,
- (i) 2-Sylow subgroup of order 4 (iii) 3-Sylow subgroup as abelian group
(ii) 5-Sylow subgroup of order 5 (iv) 3-Sylow is cyclic of 3
(v) none of these
39. $\int_0^a \int_0^b dx dy = \dots?$
- (i) a (iv) $a-b$
(ii) b (v) ab
(iii) $\frac{a}{b}$
40. $\int_0^a \int_0^b \int_0^c dx dy dz = \dots?$
- (i) $\frac{ab}{c}$ (iv) $a+b+c$
(ii) $\frac{a}{bc}$ (v) abc
(iii) $ab+c$
41. A cyclic group of order 8 has n , conjugate classes where $n = ?$
- (i) 2 (iv) 8
(ii) 4 (v) none of these
(iii) 6
42. Which of the following groups you find is cyclic ?
- (i) $C_2 \times C_4$ (iv) $C_3 \times C_4$
(ii) $C_2 \times C_6$ (v) $C_4 \times C_6$
(iii) $C_3 \times C_6$

43. Let $(R, +, \cdot)$ be a ring. Then
- | | |
|---------------------------------|--|
| (i) $(R, +)$ is a simple group | (iv) (R, \cdot) is a symmetric group |
| (ii) $(R, +)$ is a cyclic group | (v) (R, \cdot) is a semi-group |
| (iii) (R, \cdot) is a group | |
44. Let $h : G \rightarrow \bar{G}$ be a group homomorphism from group G into \bar{G} if g in G is of order 10 then
- | | |
|-----------------------------|---------------------------|
| (i) $h(g)$ is of order 3 | (iv) $h(g)$ is of order 4 |
| (ii) $h(g)$ is of order 7 | (v) none of these |
| (iii) $h(g)$ is of order 12 | |
45. Let $T : R^5 \rightarrow R^3$ be a linear transformation from R^5 into R^3 such that $\text{Ker}T$ is a subspace of R^5 of dimension 3. Then $T(R^5)$ is a
- | | |
|-------------------------------|---------------------------|
| (i) trivial subspace of R^5 | (iv) subspace of degree 3 |
| (ii) subspace of degree 1 | (v) none of these |
| (iii) subspace of dimension 2 | |
46. If $x < 0$ then $|x| = \dots\dots\dots$, for a real number x ,
- | | |
|-------------|-------------------|
| (i) $\pm x$ | (iv) 0 |
| (ii) x | (v) none of these |
| (iii) $-x$ | |
47. The inequality $\frac{x}{x-1} \geq 0$ holds true for,
- | | |
|-----------------------------|----------------------------|
| (i) $x > 1$ | (iv) $0 < x < \frac{1}{4}$ |
| (ii) $0 < x < 1$ | (v) none of these. |
| (iii) $0 < x < \frac{1}{2}$ | |
48. A topological space X is a T_1 -space iff each singleton in X is,
- | | |
|-----------------|-------------------|
| (i) open | (iv) half closed |
| (ii) closed | (v) none of these |
| (iii) half open | |
49. The topology defined on the real line is called;
- | | |
|-------------------------|---------------------|
| (i) left ray topology | (iv) usual topology |
| (ii) right ray topology | (v) none of these |
| (iii) cofinite topology | |
50. The real space R is a
- | | |
|----------------------------|-------------------|
| (i) first category space | (iv) finite Space |
| (ii) second category space | (v) open space |
| (iii) closed space | |
51. The smallest field should have,
- | | |
|----------------------|--------------------|
| (i) one element | (iv) four elements |
| (ii) two elements | (v) ten elements |
| (iii) three elements | |

(P.T.O)

52. Let $\varphi : G \rightarrow \overline{G}$ be an injective homomorphism from a group G into \overline{G} . The $\text{Ker } \varphi = \dots$
- | | |
|----------------------|-------------------|
| (i) G | (iv) φ |
| (ii) $\{e\}$ | (v) none of these |
| (iii) \overline{G} | |
53. For a ring R , a group M forms an R -module. Then
- | | |
|-----------------------------------|---------------------------------------|
| (i) M is a multiplicative group | (iii) M is a additive abelian group |
| (ii) M is necessarily cyclic | (iv) M is a simple group |
| | (v) none of these |
54. Let G be an abelian group. Then
- | | |
|-----------------------------------|------------------------------------|
| (i) the centre of $G = \{e\}$ | (iv) the centre of G is infinite |
| (ii) the centre of $G = \varphi$ | (v) the centre of $G = G$ |
| (iii) the centre of G is finite | |
55. Let G be a cyclic group of order 6. Then
- | | |
|--|--------------------------------|
| (i) G is simple group | (iii) $G \cong C_2 \times C_3$ |
| (ii) G has one non abelian sub-group | (iv) G is non-abelian |
| | (v) none of these |
56. If $X = \{a, b, c\}$ and $\tau = \{\varphi, \{x\}, \{a, b\}, X\}$ is a topology on X . Then the neighbourhood system $N(a)$ of a in X is
- | | |
|-----------------------|---|
| (i) $\{\{a, b\}, X\}$ | (iv) $N \{\{a\}, \{a, b\}, \{a, c\}, X\}$ |
| (ii) $\{X\}$ | (v) none of these |
| (iii) $\{\{a\}, X\}$ | |
57. For a, b in \mathbb{R} , the property; " $a > b$ or $a = b$ or $a < b$ " is called as;
- | | |
|-----------------------------------|-----------------------|
| (i) cancellation property | (iv) inverse property |
| (ii) left distributive property | (v) none of these |
| (iii) right distributive property | |
58. Let $G = \{1, w, w^2\}$ be the group of all cube roots of unity and $F = \{0, 1\}$ be a field. Then FG -algebra has
- | | |
|---------------------|--------------------|
| (i) two elements | (iv) nine elements |
| (ii) three elements | (v) none of these |
| (iii) four elements | |
59. The alternating group A_4 is
- | | |
|--------------|-------------------|
| (i) abelian | (iv) non-abelian |
| (ii) cyclic | (v) none of these |
| (iii) simple | |
60. The general linear group $GL_2(F_q)$ has order,
- | | |
|-----------------|-----------------------|
| (i) q^2-1 | (iv) $(q^2-1)(q^2-q)$ |
| (ii) q^2+1 | (v) none of these |
| (iii) q^2+q+1 | |

